

What is claimed is:

1	1. A method of identifying a nucleotide in at least a first position in a			
2	polynucleotide sequence, comprising:			
3	providing a polynucleotide target sequence;			
4	hybridizing the target sequence with a first oligonucleotide probe, wherein:			
5	the probe comprises a first subsequence of nucleotides, a first terminal nucleotide,			
6	and a first florescent label;			
7	the subsequence is complementary to a portion of the target sequence that is			
8	immediately adjacent to the first position; and			
9	the terminal nucleotide is complementary to one possible nucleotide in the first			
0	position;			
1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	contacting the hybridized probe and target sequence with polymerase extension			
	reagents in a first extension reaction mixture;			
	monitoring a fluorescent signal from the first extension reaction mixture that is			
	indicative of the presence or absence of polymerase extension of the probe, the presence of			
	polymerase extension of the probe indicating that the terminal nucleotide is complementary to the			
6	nucleotide in the first position; and			
7	identifying the nucleotide in the first position.			
A with the time to	2. The method of claim 1, wherein the terminal nucleotide is a 3'-terminal nucleotide.			
1	3. The method of claim 2, wherein the fluorescent label is coupled to the 3'-			
2	terminal nucleotide.			
1	4. The method of claim 1, wherein the polymerase extension reagents include			
2	3'-5' DNA polymerase enzyme.			
1	5. The method of claim 1, wherein the first fluorescent label is coupled to the			
2	first terminal nucleotide.			

1	0.	The method of claim 1, wherein the probe is from about 10 to about 50	
2	nucleotides in length.		
1 2	7. nucleotides in length.	The method of claim 1, wherein the subsequence is from about 9 to about 49	
1 2	8. a non-proofreading po	The method of claim 1, wherein the polymerase extension reagents comprise slymerase.	
1 2	9. from exonucleoase mi	The method of claim 8, wherein the non-proofreading polymerase is selected nus klenow fragment, Taq polymerase, and Thermosequanase.	
1	10. microfluidic device.	The method of claim 1, wherein the contacting step occurs in a channel of a	
	*	The method of claim 1, wherein the monitoring step comprises monitoring a rescence emitted from the first extension reaction mixture, a decrease in a indicating the presence of polymerase extension.	
	·	The method of claim 1, further comprising: zing the target sequence with a second oligonucleotide probe that comprises: the first subsequence of nucleotides, a second terminal nucleotide, and a cl, the second fluorescent label being distinguishable from the first fluorescent	
5	label; and		
6		the second terminal nucleotide is different from the first terminal nucleotide	
7	and is complementary to one possible nucleotide in the first position; and		
8	wherein the monitoring step comprises monitoring fluorescent signals from each of		
9	the first and second fluorescent labels, the fluorescent signal from one of the first and second		
10	fluorescent labels beir	ng indicative of polymerase extension of the first or second oligonucleotide	
11	probe, respectively.		

1	13. A method for identifying a nucleotide in a first position in a target fluctere			
2	acid sequence, comprising:			
3	amplifying the target nucleic acid sequence in a first reaction mixture that includes			
4	effective amounts of polymerase enzyme and four dNTPs;			
5	introducing into the first reaction mixture a first primer sequence to produce a secon			
6	reaction mixture under conditions conducive to a polymerase mediated primer extension, wherein			
7	the first primer sequence comprises a first subsequence of nucleotides, a first terminal nucleotide,			
8	and a first florescent label, wherein the subsequence is complementary to a portion of the target			
9	sequence that is immediately adjacent to the first position, and the first terminal nucleotide is			
10	complementary to one possible nucleotide in the first position;			
11	monitoring a fluorescent signal from the second reaction mixture that is indicative of			
12	a presence or absence of extension of the first primer sequence; and			
13	identifying the nucleotide in the first position based upon whether the fluorescent			
	signal is indicative of the presence of extension of the first primer sequence.			
1 I	14. A system for identifying at least a first nucleotide in a target nucleic acid			
1.2	sequence, comprising:			
11 📆	a reaction vessel having disposed therein:			
	a first target nucleic acid sequence having an unknown nucleotide at a first position;			
Ğ	a first oligonucleotide probe having a first subsequence of nucleotides, the			
7	first subsequence being complementary to a subsequence of nucleotides in the target sequence that			
8	are immediately adjacent to the first position, a first terminal nucleotide that is positioned to be			
9	adjacent to the first position when the first subsequence of the probe is hybridized to the			
10	subsequence of the target, and a first fluorescent label; and			
11	polymerase extension reagents; and			
12	a detector configured to monitor a fluorescent signal from the reaction vessel that is			
13	indicative of a presence or absence of polymerase extension of the probe.			

The system of claim 14, wherein the first fluorescent label is coupled to the

1

2

15.

first terminal nucleotide.

- 1 16. The system of claim 14, wherein the first terminal nucleotide is a 3' terminal nucleotide, and the polymerase extension reagents include a 3'-5' DNA polymerase enzyme.
- 1 The system of claim 14, wherein the reaction vessel is selected from a 2 reaction well in a multiwell plate, a capillary channel and a channel in a microfluidic channel 3 network.
- 1 18. The system of claim 14, wherein the reaction vessel further comprises a 2 second oligonucleotide probe disposed therein, the second oligonucleotide probe comprising the 3 first subsequence of nucleotides, a second terminal nucleotide different from the first terminal 4 nucleotide, that is complementary to one possible nucleotide in the first position, and a second 5 fluorescent label.

The training that the training that the